OFFICIAL

SAPPG Upload to SA Health Website and Practice Guidelines Web-Based App Summary

Date uploaded: 17/04/2024

Title of Document (previous title if applicable)	Revised or New	Summary of key points / changes	Risk management concerns +/- budget implications
 Postpartum Haemorrhage Note that the following PPGs have been merged into the PPH PPG and removed from circulation: Ergot derivatives Prostaglandin analogues Secondary PPH Intra-uterine balloon tamponade Oxytocin prophylaxis for third stage management and PPH 	Revised (version 6)	 This PPG has been reviewed in line with current international and national guidelines, and latest research recommendations for the management of PPH. Key points: Use of Carbetocin for active management of third stage of labour has been added. To be used in vaginal births only when women meet the criteria of: PPH risk factors and contraindications to syntrometrine/ergometrine and/or presence of hypertension/cardiac anomalies. Note: Carbetocin is not currently on the SA formulary for vaginal births therefore an IPU form must be completed and sent to pharmacy. Addition of tranexamic acid as part of first line PPH management medication Addition of Carboprost as part of second line PPH management medication New primary PPH management algorithm that includes tranexamic acid and carboprost. External link for larger, printable flowchart – access via the link at the bottom of page 2 of the PPG. Print as needed. Secondary PPH management flowchart updated with improved readability. All blood loss must be measured (e.g., weighing soiled linen/pads). Previous recommendation for visual estimation no longer applicable as not evidence based. Educate all staff on how to measure and record blood loss using appropriate measuring aids. Previous odds ratio risk tables removed as supported by low quality evidence. Risk factors for PPH table updated. PPH record form updated to reflect new PPH management flowchart. Suggested citation added (New) Appendices updated. 	Distribution of information to all clinicians across SA Health, via notification of upload. All relevant LHNs to review local guidelines on PPH management and adjust accordingly. This PPG has been presented to the perinatal emergency education strategy (PEES) group. Ensure all staff involved in management of PPH are aware of practice changes.